

DYNAMICS OF A BEC IN THE THOMAS-FERMI REGIME

Daniele Dimonte

with E. L. Giacomelli, M. Correggi and P. Pickl
SwissMAP General Meeting, 08.09.2020



**University
of Basel**

TABLE OF CONTENT

INTRODUCTION

DYNAMICAL PROBLEM

CONCLUSION

INTRODUCTION

DYNAMICAL PROBLEM

CONCLUSION

PHYSICAL SETTING

Goal: study the dynamics of N identical bosons in a box Λ with periodic BC

- **Thermodynamic limit:** with fixed density $\rho := N/|\Lambda|$, study of the limit of infinite volume of the **energy per particle**

$$\begin{aligned} \epsilon(\rho) &:= \lim_{N \rightarrow +\infty} \frac{\inf \sigma(H_N)}{N} \\ &= 4\pi\rho a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} + o\left(\sqrt{\rho a^3}\right) \right) \quad (\text{LHY}) \end{aligned}$$

- Dilute limit: if ρa^3 is small (a scattering length, effective length of the interaction) we obtain the Lee-Huang-Yang formula (LHY)

PHYSICAL SETTING

Goal: study the dynamics of N identical bosons in a box Λ with periodic BC

- **Thermodynamic limit:** with fixed density $\rho := N/|\Lambda|$, study of the limit of infinite volume of the **energy per particle**

$$\begin{aligned} \epsilon(\rho) &:= \lim_{N \rightarrow +\infty} \frac{\inf \sigma(H_N)}{N} \\ &= 4\pi\rho a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} + o\left(\sqrt{\rho a^3}\right) \right) \quad (\text{LHY}) \end{aligned}$$

- **Dilute limit:** if ρa^3 is small (a scattering length, effective length of the interaction) we obtain the Lee-Huang-Yang formula (LHY)

PHYSICAL SETTING

For the variational problem, in a dilute limit (at $T = 0$) one expects that the macroscopic ground state of the system Ψ^{GS} is well approximated by a one-particle state, i.e., there is **Bose-Einstein Condensation (BEC)**

$$H_N \Psi^{\text{GS}} = E_0(N) \Psi^{\text{GS}}$$
$$\Psi^{\text{GS}} \approx (\varphi^{\text{GS}})^{\otimes N}$$

φ^{GS} ground state of a nonlinear effective one-particle functional

$$\mathcal{E}^{\text{eff}}[\varphi] := \langle \varphi, h\varphi \rangle + \langle \varphi, \mathcal{V}_{\text{eff}}(\varphi) \rangle$$

with h one-particle Hamiltonian and \mathcal{V}_{eff} an effective nonlinear potential

DILUTE LIMITS

Let v_N be the (N -dependent) pair interaction

- Mean-Field (Hartree)

$$v_N(\mathbf{x}) := \frac{1}{N} v(\mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} \left(v * |\psi|^2 \right) |\psi|^2$$

- Gross-Pitaevskii (GP)

$$v_N(\mathbf{x}) := N^2 v(N\mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} g |\psi|^4$$

- Intermediate regimes ($\beta \in (0, 1)$)

$$v_N(\mathbf{x}) := N^{3\beta-1} v(N^\beta \mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} \left(\int v \right) |\psi|^4$$

In all these cases a_N the scattering length of v_N satisfies $8\pi N a_N \rightarrow g$, with g constant ($\rho a_N^3 \approx N^{-2} \ll 1$)

DILUTE LIMITS

Let v_N be the (N -dependent) pair interaction

- Mean-Field (Hartree) ($\beta = 0$)

$$v_N(\mathbf{x}) := \frac{1}{N} v(\mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} \left(v * |\psi|^2 \right) |\psi|^2$$

- Gross-Pitaevskii (GP) ($\beta = 1$)

$$v_N(\mathbf{x}) := N^2 v(N\mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} g |\psi|^4$$

- Intermediate regimes ($\beta \in (0, 1)$)

$$v_N(\mathbf{x}) := N^{3\beta-1} v(N^\beta \mathbf{x}), \quad \mathcal{V}_{\text{eff}}(\psi) = \frac{1}{2} \left(\int v \right) |\psi|^4$$

In all these cases a_N the scattering length of v_N satisfies $8\pi N a_N \rightarrow g$, with g constant ($\rho a_N^3 \approx N^{-2} \ll 1$)

THOMAS-FERMI REGIME

In experimental settings, in particular in considering rotating systems, $Na_N \gg 1$; this is called **Thomas-Fermi** regime, in analogy with the density theory for large atoms

We consider a pair interaction such that $8\pi a_N \rightarrow +\infty$, compatibly with the dilute condition $\rho a_N^3 \ll 1$

THOMAS-FERMI REGIME

Fix the size of Λ and consider the following many-body Hamiltonian

$$H_N := \sum_{j=1}^N (-\Delta_j) + g_N N^{3\beta-1} \sum_{1 \leq j < k \leq N} v(N^\beta (\mathbf{x}_j - \mathbf{x}_k))$$

defined on $\mathcal{H}_N := \mathfrak{h}^{\otimes_s N}$, with $\mathfrak{h} = L^2(\Lambda)$

- Without loss of generality $\int v = 1$; then the scattering length of $g_N N^{3\beta-1} v(N^\beta \cdot)$ is given for $\beta \in [0, 1)$ by

$$Na_N = \frac{1}{8\pi} g_N (1 + o(1))$$

therefore we require $g_N \gg 1$ (TF regime)

- If $g_N \leq N^{2/3}$ this is still a *dilute limit*

MATHEMATICAL SETTING

To evaluate one-particle observables on many-body states $\Psi \in \mathcal{H}_N$ it is convenient to introduce the **1-particle reduced density matrix** $\gamma_\Psi^{(1)}$ defined so that

$$\left\langle \Psi, \sum_{j=1}^N A_j \Psi \right\rangle = N \operatorname{tr} \left[\gamma_\Psi^{(1)} A \right]$$

for any A a one-particle observable

COMPLETE BEC

Given a many-body state $\Psi \in \mathcal{H}_N$ and a one-particle state $\varphi \in \mathfrak{h}$

$$\gamma_\Psi^{(1)} \rightarrow P_\varphi := |\varphi\rangle \langle \varphi|, \quad \text{in } \mathfrak{S}_1(\mathfrak{h})$$

i.e., a macroscopic fraction of the particles occupies the same one-particle state

INTRODUCTION

DYNAMICAL PROBLEM

CONCLUSION

SETTING

We consider a trapped system in $\Lambda = [-\frac{1}{2}, \frac{1}{2}]^3$ ($\hbar = L^2(\Lambda)$)

$$H_N := \sum_{j=1}^N (-\Delta_j) + g_N N^{3\beta-1} \sum_{1 \leq j < k \leq N} v(N^\beta(\mathbf{x}_j - \mathbf{x}_k))$$

The solution to the Schrödinger equation is

$$\begin{cases} i\partial_t \Psi_{N,t} = H_N \Psi_{N,t} \\ \Psi_{N,t}|_{t=0} = \Psi_{N,0} \end{cases}$$

Goal: understand whether complete BEC is preserved by time evolution, i.e.

$$\gamma_{\Psi_{N,0}}^{(1)} \rightarrow P_{\varphi_0} \text{ in } \mathfrak{S}_1(\hbar) \implies \gamma_{\Psi_{N,t}}^{(1)} \rightarrow P_{\varphi_t^{\text{GP}}} \text{ in } \mathfrak{S}_1(\hbar)$$

GROSS-PITAEVSKII EQUATION

Expected limiting equation: the **time-dependent GP equation**

$$\begin{cases} i\partial_t \varphi_t^{\text{GP}} = -\Delta \varphi_t^{\text{GP}} + g_N |\varphi_t^{\text{GP}}|^2 \varphi_t^{\text{GP}} \\ \varphi_t^{\text{GP}}|_{t=0} = \varphi_0 \end{cases}$$

Energy of the system:

$$\begin{aligned} \mathcal{E}^{\text{GP}}[\varphi] &= \int_{\Lambda} d\mathbf{x} \left(\frac{1}{2} |\nabla \varphi(\mathbf{x})|^2 + \frac{g_N}{2} |\varphi(\mathbf{x})|^4 \right) \\ E^{\text{GP}} &= \inf_{\|\varphi\|_2=1} \mathcal{E}^{\text{GP}}[\varphi] \end{aligned}$$

Idea: for low energies the kinetic term is negligible if N is large

THOMAS-FERMI ENERGY

Dropping the kinetic term we obtain the TF energy functional

$$\mathcal{E}^{\text{TF}}[\rho] = \frac{gN}{2} \int_{\Lambda} d\mathbf{x} \rho^2(\mathbf{x}),$$
$$E^{\text{TF}} = \inf_{\|\rho\|_1=1, \rho \geq 0} \mathcal{E}^{\text{TF}}[\rho]$$

Fact: in a box $E^{\text{GP}} = E^{\text{TF}} = \frac{gN}{2}$
(in \mathbb{R}^3 , $E^{\text{GP}} \approx E^{\text{TF}}$ at first order in N)

INTERMEDIATE EQUATION

To prove the approximation $\gamma_{\Psi_{N,t}}^{(1)} \approx P_{\varphi_t^{\text{GP}}}$ it is helpful to introduce an intermediate effective equation, the **time-dependent Hartree (H) equation**

$$\begin{cases} i\partial_t \varphi_t^{\text{H}} = -\Delta \varphi_t^{\text{H}} + g_N v_N * |\varphi_t^{\text{H}}|^2 \varphi_t^{\text{H}} \\ \varphi_t^{\text{H}}|_{t=0} = \varphi_0 \end{cases}$$

We exploit $v_N * |\varphi|^2 \rightarrow |\varphi|^2$, but we need control on $\|\varphi\|_\infty$ independent on g_N

INTERMEDIATE EQUATION

To prove the approximation $\gamma_{\Psi_{N,t}}^{(1)} \approx P_{\varphi_t^{\text{GP}}}$ it is helpful to introduce an intermediate effective equation, the **time-dependent Hartree (H) equation**

$$\begin{cases} i\partial_t \varphi_t^{\text{H}} = -\Delta \varphi_t^{\text{H}} + g_N v_N * |\varphi_t^{\text{H}}|^2 \varphi_t^{\text{H}} \\ \varphi_t^{\text{H}}|_{t=0} = \varphi_0 \end{cases}$$

We exploit $v_N * |\varphi|^2 \rightarrow |\varphi|^2$, but we need control on $\|\varphi\|_\infty$
independent on g_N

CONJECTURE

Let φ_0 be the initial datum of the GP equation

$$\varphi_0 \in L^\infty(\Lambda) \implies \sup_{t \in \mathbb{R}} \|\varphi_t^H\|_\infty \leq C$$

THEOREM

Assume that $v \in L^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3, dx)$, the Conjecture holds true and

$$\begin{aligned} \left\| \gamma_{\Psi_{N,0}}^{(1)} - P_{\varphi_0^{\text{GP}}} \right\|_{\mathfrak{S}^1} &\ll N^{-\frac{1-3\beta}{2}} \\ \mathcal{E}^{\text{GP}}[\varphi_0] - E^{\text{GP}} &\ll \xi_N \leq \sqrt{g_N} \\ g_N &\ll \log N \end{aligned}$$

then for each $t \in \mathbb{R}$ and for any $\beta \in [0, 1/6)$ there is *complete BEC* on φ_t^{GP} , i.e.

$$\left\| \gamma_{\Psi_{N,t}}^{(1)} - P_{\varphi_t^{\text{GP}}} \right\|_{\mathfrak{S}^1} \ll 1$$

REMARKS

- Similar result is achievable also in $d = 2$
- Open question is to go beyond $\beta = 1/6$; also related to stationary problem limitations
- (HP1) means that there is BEC in the initial datum $\Psi_{N,0}$ on the state φ_0
- (HP2) means that the GP initial datum φ_0 is close to a ground state in energy: important to prove that the Hartree solution is close to the GP solution
- (HP3) is necessary to prove condensation on a state φ_t^H ; still allows for a dilute limit

$$\left\| \gamma_{\Psi_{N,0}}^{(1)} - P_{\varphi_0^{\text{GP}}} \right\|_{\mathfrak{S}^1} \ll N^{-\frac{1-3\beta}{2}} \quad (\text{HP1})$$

$$\mathcal{E}^{\text{GP}}[\varphi_0] - E^{\text{GP}} \ll \xi_N \leq \sqrt{g_N} \quad (\text{HP2})$$

$$g_N \ll \log N \quad (\text{HP3})$$

SKETCH OF THE PROOF

Two parts:

- Approximate the $\gamma_{\Psi_{N,t}}^{(1)}$ with $P_{\varphi_t^H}$
- Estimate the difference between φ_t^H and φ_t^{GP}

Main ingredients:

- Tools developed in [P11]
- Energy estimates for the one-particle problem

MANY-BODY TO HARTREE

Similarly as in **[P11]**, the goal is obtaining a Grönwall-type estimate for

$$\alpha_t := 1 - \langle \Psi_{N,t}, (|\varphi_t^H\rangle \langle \varphi_t^H|)_1 \Psi_{N,t} \rangle$$

We need to estimate terms of the form

$$\left\| v_N * |\varphi_t^H|^2 \right\|_\infty \leq \|v\|_1 \|\varphi_t^H\|_\infty^2$$

Using the Conjecture we get the desired result; if we do not assume it, we can only use the kinetic energy: *we do not reach the time scale of vortices* (compare with **[JS15]**)

[JS15] Jerrard, Smets, “Vortex dynamics for the two-dimensional non-homogeneous Gross-Pitaevskii equation”

HARTREE TO GROSS-PITAEVSKII

$$\begin{aligned}
\partial_t \|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_2^2 &\leq gN \left| \text{Im} \langle \varphi_t^{\text{H}}, \left(|\varphi_t^{\text{GP}}|^2 - v_N * |\varphi_t^{\text{H}}|^2 \right) \varphi_t^{\text{GP}} \rangle \right| \\
&\leq gN \left| \langle \varphi_t^{\text{H}}, \left(|\varphi_t^{\text{GP}}|^2 - |\varphi_t^{\text{H}}|^2 \right) \varphi_t^{\text{GP}} \rangle \right| \\
&\quad + gN \left| \langle \varphi_t^{\text{H}}, \left(|\varphi_t^{\text{H}}|^2 - v_N * |\varphi_t^{\text{H}}|^2 \right) \varphi_t^{\text{GP}} \rangle \right|
\end{aligned}$$

To prove convergence of this last two terms use L^2 difference of the square of the solutions (energy bound) for the first term and $v_N \rightarrow \delta$ as a distribution for the second one:

$$\begin{aligned}
&\left| \langle \varphi_t^{\text{H}}, \left(|\varphi_t^{\text{H}}|^2 - v_N * |\varphi_t^{\text{H}}|^2 \right) \varphi_t^{\text{GP}} \rangle \right| \leq \\
&\leq \frac{C}{N^\beta} \|\nabla \varphi_t^{\text{H}}\|_2 \|\varphi_t^{\text{H}}\|_\infty \|\varphi_t^{\text{H}}\|_4 \|\varphi_t^{\text{GP}}\|_4
\end{aligned}$$

INTRODUCTION

DYNAMICAL PROBLEM

CONCLUSION

CONCLUSION

- Condensation is preserved under suitable assumptions of regularity on the solution
 - Q: How to prove the Conjecture?
 - Q: Vortices are encoded in the vorticity measure, which depends on the gradient of the solution; can a similar result be proven in a stronger (e.g. H^1) norm?
- There is BEC in the Thomas Fermi limit, at least in a scaling with $\beta < 1/3$ (work in progress with M. Correggi and E. L. Giacomelli)
 - Q: Can we extend the result for $\beta > 1/6$?

CONCLUSION

- Condensation is preserved under suitable assumptions of regularity on the solution
 - Q: How to prove the Conjecture?
 - Q: Vortices are encoded in the vorticity measure, which depends on the gradient of the solution; can a similar result be proven in a stronger (e.g. H^1) norm?
- There is BEC in the Thomas Fermi limit, at least in a scaling with $\beta < 1/3$ (work in progress with M. Correggi and E. L. Giacomelli)
 - Q: Can we extend the result for $\beta > 1/6$?

Thanks for the attention!